

ASYMPTOTIC ESTIMATES FOR PHI FUNCTIONS FOR SUBSETS OF $\{m+1, m+2, \dots, n\}$

MELVYN B. NATHANSON AND BROOKE OROSZ

ABSTRACT. Let $f(m, n)$ denote the number of relatively prime subsets of $\{m+1, m+2, \dots, n\}$, and let $\Phi(m, n)$ denote the number of subsets A of $\{m+1, m+2, \dots, n\}$ such that $\gcd(A)$ is relatively prime to n . Let $f_k(m, n)$ and $\Phi_k(m, n)$ be the analogous counting functions restricted to sets of cardinality k . Simple explicit formulas and asymptotic estimates are obtained for these four functions.

A nonempty set A of integers is called *relatively prime* if $\gcd(A) = 1$. Let $f(n)$ denote the number of nonempty relatively prime subsets of $\{1, 2, \dots, n\}$ and, for $k \geq 1$, let $f_k(n)$ denote the number of relatively prime subsets of $\{1, 2, \dots, n\}$ of cardinality k .

Euler's phi function $\varphi(n)$ counts the number of positive integers a in the set $\{1, 2, \dots, n\}$ such that a is relatively prime to n . The Phi function $\Phi(n)$ counts the number of nonempty subsets A of the set $\{1, \dots, n\}$ such that $\gcd(A)$ is relatively prime to n or, equivalently, such that $A \cup \{n\}$ is relatively prime. For every positive integer k , the function $\Phi_k(n)$ counts the number of sets $A \subseteq \{1, \dots, n\}$ such that $\text{card}(A) = k$ and $\gcd(A)$ is relatively prime to n .

Nathanson [2] introduced these four functions for subsets of $\{1, 2, \dots, n\}$, and El Bachraoui [1] generalized them to subsets of the set $\{m+1, m+2, \dots, n\}$ for arbitrary nonnegative integers $m < n$.¹ We shall obtain simple explicit formulas and asymptotic estimates for the four functions.

For every real number x , we denote by $[x]$ the greatest integer not exceeding x . We often use the elementary inequality $[x] - [y] \leq [x - y] + 1$ for all $x, y \in \mathbf{R}$.

Theorem 1. *For nonnegative integers $m < n$, let $f(m, n)$ denote the number of relatively prime subsets of $\{m+1, m+2, \dots, n\}$. Then*

$$f(m, n) = \sum_{d=1}^n \mu(d) \left(2^{\lfloor n/d \rfloor - \lfloor m/d \rfloor} - 1 \right)$$

and

$$0 \leq 2^{n-m} - 2^{\lfloor n/2 \rfloor - \lfloor m/2 \rfloor} - f(m, n) \leq 2n2^{\lfloor (n-m)/3 \rfloor}.$$

Date: February 1, 2008.

2000 Mathematics Subject Classification. Primary 11A25, 11B05, 11B13, 11B75.

Key words and phrases. Relatively prime sets, Euler's phi function, Nathanson's phi function, combinatorial number theory, elementary number theory.

The work of M.B.N. was supported in part by grants from the NSA Mathematical Sciences Program and the PSC-CUNY Research Award Program.

¹Actually, our function $f(m, n)$ is El Bachraoui's function $f(m+1, n)$, and similarly for the other three functions. This small change yields formulas that are more symmetric and pleasing esthetically.

Proof. El Bachraoui [1] proved that

$$f(m, n) = \sum_{d=1}^n \mu(d) \left(2^{\lfloor n/d \rfloor} - 1 \right) - \sum_{i=1}^m \sum_{d|i} \mu(d) 2^{\lfloor n/d \rfloor - i/d}.$$

Rearranging this identity, we obtain

$$\begin{aligned} f(m, n) &= \sum_{d=1}^n \mu(d) \left(2^{\lfloor n/d \rfloor} - 1 \right) - \sum_{d=1}^m \mu(d) 2^{\lfloor n/d \rfloor} \sum_{\substack{i=1 \\ i|d}}^m 2^{-i/d} \\ &= \sum_{d=1}^n \mu(d) \left(2^{\lfloor n/d \rfloor} - 1 \right) - \sum_{d=1}^m \mu(d) 2^{\lfloor n/d \rfloor} \sum_{j=1}^{\lfloor m/d \rfloor} 2^{-j} \\ &= \sum_{d=1}^n \mu(d) 2^{\lfloor n/d \rfloor} \left(1 - \sum_{j=1}^{\lfloor m/d \rfloor} 2^{-j} \right) - \sum_{d=1}^n \mu(d) \\ &= \sum_{d=1}^n \mu(d) \left(2^{\lfloor n/d \rfloor - \lfloor m/d \rfloor} - 1 \right). \end{aligned}$$

Let $d \in \{1, 2, \dots, n\}$. Then $m+1 \leq a \leq n$ and d divides a if and only if $\lfloor m/d \rfloor + 1 \leq a/d \leq \lfloor n/d \rfloor$. It follows that $A \subseteq \{m+1, \dots, n\}$ and $\gcd(A) = d$ if and only if $A' = (1/d) * A \subseteq \{\lfloor m/d \rfloor + 1, \dots, \lfloor n/d \rfloor\}$ and $\gcd(A') = 1$. Therefore,

$$\begin{aligned} 2^{n-m} - 1 &= \sum_{d=1}^n f(\lfloor m/d \rfloor, \lfloor n/d \rfloor) \\ &\leq f(m, n) + 2^{\lfloor n/2 \rfloor - \lfloor m/2 \rfloor} - 1 + \sum_{d=3}^n 2^{\lfloor n/d \rfloor - \lfloor m/d \rfloor} \end{aligned}$$

and we obtain the lower bound

$$f(m, n) \geq 2^{n-m} - 2^{\lfloor n/2 \rfloor - \lfloor m/2 \rfloor} - 2n2^{[(n-m)/3]}.$$

For the upper bound, we observe that the number of subsets of even integers contained in the set $\{m+1, \dots, n\}$ is exactly $2^{\lfloor n/2 \rfloor - \lfloor m/2 \rfloor}$ and so

$$f(m, n) \leq 2^{n-m} - 2^{\lfloor n/2 \rfloor - \lfloor m/2 \rfloor}.$$

This completes the proof. \square

Theorem 2. For nonnegative integers $m < n$ and for $k \geq 1$, let $f_k(m, n)$ denote the number of relatively prime subsets of $\{m+1, m+2, \dots, n\}$ of cardinality k . Then

$$f_k(m, n) = \sum_{d=1}^n \mu(d) \binom{\lfloor n/d \rfloor - \lfloor m/d \rfloor}{k}$$

and

$$0 \leq \binom{n-m}{k} - \binom{\lfloor n/2 \rfloor - \lfloor m/2 \rfloor}{k} - f_k(m, n) \leq n \binom{[(n-m)/3] + 2}{k}.$$

Proof. El Bachraoui [1] proved that

$$f_k(m, n) = \sum_{d=1}^n \mu(d) \binom{\lfloor n/d \rfloor}{k} - \sum_{i=1}^m \sum_{d|i} \mu(d) \binom{\lfloor n/d \rfloor - i/d}{k-1}.$$

We recall the combinatorial fact that for $k \geq 1$ and $0 \leq M \leq N$, we have

$$\binom{N}{k} - \sum_{j=1}^M \binom{N-j}{k-1} = \binom{N-M}{k}.$$

Then

$$\begin{aligned} f_k(m, n) &= \sum_{d=1}^n \mu(d) \binom{\lfloor n/d \rfloor}{k} - \sum_{d=1}^m \mu(d) \sum_{\substack{i=1 \\ d|i}}^m \binom{\lfloor n/d \rfloor - i/d}{k-1} \\ &= \sum_{d=1}^m \mu(d) \left(\binom{\lfloor n/d \rfloor}{k} - \sum_{j=1}^{\lfloor m/d \rfloor} \binom{\lfloor n/d \rfloor - j}{k-1} \right) + \sum_{d=m+1}^n \mu(d) \binom{\lfloor n/d \rfloor}{k} \\ &= \sum_{d=1}^m \mu(d) \binom{\lfloor n/d \rfloor - \lfloor m/d \rfloor}{k} + \sum_{d=m+1}^n \mu(d) \binom{\lfloor n/d \rfloor}{k} \\ &= \sum_{d=1}^n \mu(d) \binom{\lfloor n/d \rfloor - \lfloor m/d \rfloor}{k}. \end{aligned}$$

We obtain an upper bound for $f_k(m, n)$ by deleting k -element sets of even integers:

$$f_k(m, n) \leq \binom{n-m}{k} - \binom{\lfloor n/2 \rfloor - \lfloor m/2 \rfloor}{k}$$

and we obtain a lower bound from the identity

$$\begin{aligned} \binom{n-m}{k} &= \sum_{d=1}^n f_k(\lfloor m/d \rfloor, \lfloor n/d \rfloor) \\ &\leq f_k(m, n) + \binom{\lfloor n/2 \rfloor - \lfloor m/2 \rfloor}{k} + \sum_{d=3}^n \binom{\lfloor n/d \rfloor - \lfloor m/d \rfloor}{k} \\ &\leq f_k(m, n) + \binom{\lfloor n/2 \rfloor - \lfloor m/2 \rfloor}{k} + n \binom{\lfloor (n-m)/3 \rfloor}{k}. \end{aligned}$$

□

Theorem 3. For nonnegative integers $m < n$, let $\Phi(m, n)$ denote the number of subsets of $[m+1, n]$ such that $\gcd(A)$ is relatively prime to n . Then

$$\Phi(m, n) = \sum_{d|n} \mu(d) 2^{(n/d) - \lfloor m/d \rfloor}.$$

If p^* is the smallest prime divisor of n , then

$$0 \leq 2^{n-m} - 2^{(n/p^*) - \lfloor m/p^* \rfloor} - \Phi(m, n) \leq 2n 2^{[(n-m)/(p^*+1)]}.$$

Proof. El Bachraoui [1] proved that

$$\Phi(m, n) = \sum_{d|n} \mu(d) 2^{n/d} - \sum_{i=1}^m \sum_{d|(i, n)} \mu(d) 2^{(n-i)/d}$$

Rearranging this identity, we obtain

$$\begin{aligned}
\Phi(m, n) &= \sum_{d|n} \mu(d) 2^{n/d} - \sum_{d|n} \mu(d) \sum_{\substack{i=1 \\ d|i}}^m 2^{(n-i)/d} \\
&= \sum_{d|n} \mu(d) 2^{n/d} - \sum_{d|n} \mu(d) \sum_{j=1}^{\lfloor m/d \rfloor} 2^{(n-jd)/d} \\
&= \sum_{d|n} \mu(d) 2^{n/d} \left[1 - \sum_{j=1}^{\lfloor m/d \rfloor} 2^{-j} \right] \\
&= \sum_{d|n} \mu(d) 2^{(n/d) - \lfloor m/d \rfloor}.
\end{aligned}$$

Let p^* be the smallest prime divisor of n . Deleting all subsets of $\{m+1, \dots, n\}$ whose elements are all multiples of p^* , we obtain the upper bound

$$\Phi(m, n) \leq 2^{n-m} - 2^{(n/p^*) - \lfloor m/p^* \rfloor}.$$

For the lower bound, we have

$$\begin{aligned}
\Phi(m, n) - \left(2^{n-m} - 2^{(n/p^*) - \lfloor m/p^* \rfloor} \right) &= \sum_{\substack{d|n \\ d > p^*}} \mu(d) 2^{(n/d) - \lfloor m/d \rfloor} \\
&\leq 2 \sum_{\substack{d|n \\ d > p^*}} 2^{[(n-m)/d]} \leq 2n 2^{[(n-m)/(p^*+1)]}.
\end{aligned}$$

This completes the proof. \square

Theorem 4. *For nonnegative integers $m < n$, let $\Phi_k(m, n)$ denote the number of subsets of cardinality k contained in the interval of integers $\{m+1, m+2, \dots, n\}$ such that $\gcd(A)$ is relatively prime to n . Then*

$$\Phi_k(m, n) = \sum_{d|n} \mu(d) \binom{n/d - \lfloor m/d \rfloor}{k}$$

and

$$0 \leq \binom{n-m}{k} - \binom{n/p^* - \lfloor m/p^* \rfloor}{k} - \Phi_k(m, n) \leq n \binom{[(n-m)/(p^*+1)] + 1}{k}.$$

Proof. Let p^* be the smallest prime divisor of n . El Bachraoui [1] proved that

$$\Phi_k(m, n) = \sum_{d|n} \mu(d) \binom{n/d}{k} - \sum_{i=1}^m \sum_{d|\gcd(i, n)} \mu(d) \binom{(n-i)/d}{k-1}.$$

Rearranging this identity, we obtain

$$\begin{aligned}
\Phi_k(m, n) &= \sum_{d|n} \mu(d) \binom{n/d}{k} - \sum_{d|n} \mu(d) \sum_{\substack{i=1 \\ i \nmid d}}^m \binom{(n-i)/d}{k-1} \\
&= \sum_{d|n} \mu(d) \left(\binom{n/d}{k} - \sum_{j=1}^{\lfloor m/d \rfloor} \binom{n/d-j}{k-1} \right) \\
&= \sum_{d|n} \mu(d) \binom{n/d - \lfloor m/d \rfloor}{k} \\
&\geq \binom{n-m}{k} - \binom{n/p^* - \lfloor m/p^* \rfloor}{k} - \sum_{\substack{d|n \\ d > p^*}} \binom{n/d - \lfloor m/d \rfloor}{k} \\
&\geq \binom{n-m}{k} - \binom{n/p^* - \lfloor m/p^* \rfloor}{k} - \sum_{\substack{d|n \\ d > p^*}} \binom{\lfloor (n-m)/d \rfloor + 1}{k} \\
&\geq \binom{n-m}{k} - \binom{n/p^* - \lfloor m/p^* \rfloor}{k} - n \binom{\lfloor (n-m)/(p^*+1) \rfloor + 1}{k}.
\end{aligned}$$

Deleting k -element subsets of $\{m+1, \dots, n\}$ whose elements are multiples of p^* , we get the upper bound

$$\Phi_k(m, n) \leq \binom{n-m}{k} - \binom{\lfloor n/p^* \rfloor - \lfloor m/p^* \rfloor}{k}.$$

This completes the proof. \square

REFERENCES

- [1] M. El Bachraoui, *The number of relatively prime subsets and phi functions for $\{m, m+1, \dots, n\}$* , *Integers* **7** (2007), to appear.
- [2] M. B. Nathanson, *Affine invariants, relatively prime sets, and a phi function for subsets of $\{1, 2, \dots, n\}$* , *Integers* **7** (2007), A1, 7 pp. (electronic).

DEPARTMENT OF MATHEMATICS, LEHMAN COLLEGE (CUNY), BRONX, NEW YORK 10468, AND
 SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NJ 08540
E-mail address: `melvyn.nathanson@lehman.cuny.edu`, `melvyn@ias.edu`

DEPARTMENT OF MATHEMATICS, CUNY GRADUATE CENTER, NEW YORK, NEW YORK 10036
E-mail address: `borosz@gc.cuny.edu`